

TADALAFIL (Adcirca[®], Cialis[®])

What is tadalafil (Adcirca[®], Cialis[®]) used for?

Tadalafil (Adcirca[®], Cialis[®]) is a prescription medicine used to treat pulmonary arterial hypertension (PAH). PAH is defined by high blood pressure in the lungs resulting from the pulmonary arteries—the blood vessels that carry blood from the heart to the lungs—becoming narrowed or scarred.

How does it work?

Tadalafil (Adcirca[®], Cialis[®]) is a phosphodiesterase 5 inhibitor (PDE-5 inhibitor) that lowers blood pressure in the lungs, making it easier for the heart to pump more effectively. Tadalafil (Adcirca[®], Cialis[®]) prevents PDE-5 enzymes from degrading nitric oxide in the pulmonary arteries, which helps dilate the blood vessels. Taking this medication may improve your ability to exercise and slow disease progression.

What do I need to know?

Contraindications:

Do not use tadalafil (Adcirca[®], Cialis[®]) if you:

- Have ever had an allergic reaction to tadalafil (Adcirca[®], Cialis[®]) or any of its ingredients
- Have had a prior episode of non-arteritic anterior ischemic optic neuropathy (NAION)
- Are taking medications that contain nitrates in any form (oral, spray, sublingual, ointment, and/or patch)
- Are taking riociguat (Adempas[®])

Tadalafil (Adcirca[®], Cialis[®]) should be avoided in severe liver or kidney disease.

Important information:

Avoid grapefruit juice.

If you are pregnant or may become pregnant, contact your medical team right away.

Drug interactions

The following drugs may interact with tadalafil (Adcirca[®], Cialis[®]). Please note that this is not a comprehensive list of drug interactions.

Please speak to your medical team before starting any new medication.

Using tadalafil (Adcirca[®], Cialis[®]) with blood pressure medications can lead to low blood pressure in some patients.

Medications for the treatment of fungal infections — e.g. ketoconazole (Apo-Ketoconazole[®]), itraconazole (Sporanox[®]), and voriconazole (Vfend[®])

Nitrates in any form — e.g. nitroglycerin spray, tablets, patch, or ointments, isosorbide dinitrate or mononitrate

Medications for the treatment of HIV infection — e.g. darunavir (Prezista[®]), ritonavir (Norvir[®]), nelfinavir (Viracept[®]), and saquinavir (Invirase[®])

Antibiotics — e.g. erythromycin (Eryc[®]), clarithromycin (Biaxin[®]), telithromycin (Ketek[®]), and rifampin (Rofact[®])

Medications for seizure disorders — e.g. phenytoin (Dilantin[®]), and carbamazepine (Tegretol[®])

Medications used to treat pulmonary hypertension and/or erectile dysfunction — e.g. riociguat (Adempas[®]), sildenafil (Revatio[®], Viagra[®]), and vardenafil (Levitra[®])

Please Note:

Tell your medical team about ALL the medicines you are taking including over-the-counter drugs, vitamins, minerals, natural supplements, and alternative medications.

TADALAFIL (Adcirca[®], Cialis[®])

How should I take tadalafil (Adcirca[®], Cialis[®])?

Take two tablets once a day, with or without food.

Missed dose:

If you miss a dose of tadalafil (Adcirca[®], Cialis[®]), take your tablet as soon as you remember. Do not take two doses at the same time. If it is almost time for your next dose, skip the missed dose; just take the next dose at your regular time.

What are common side effects?

Some patients may experience side effects when taking this medication. If the side effects are severe or bothersome, please contact your medical team. Side effects may include:

- Headache
- Flushing
- Muscle aches
- Nasal congestion
- Indigestion and heartburn
- Blurred or impaired vision
- Sudden vision or hearing loss

When should I notify my PH clinic or seek medical attention?

Contact your medical team or seek medical attention immediately if you develop:

- Symptoms of worsening PH, including: difficulty breathing, fluid retention or weight gain, and/or dizziness or passing out
- Sudden vision or hearing loss
- Painful or prolonged erection
- An allergic reaction, including: swelling of the mouth, tongue, face, and throat; itching; rash; and/or difficulty breathing

If you take more than your prescribed dose, contact your medical team immediately, even if you feel well.

Do not stop taking tadalafil (Adcirca[®], Cialis[®]) unless advised by your medical team.

Additional information:

Store out of reach of children at 15°–30° C.

Do not use expired medication.

Do not throw away any medication via wastewater or household waste. Ask your pharmacist how to dispose of medications you no longer require.

For more information please see full product monograph.



PULMONARY HYPERTENSION
ASSOCIATION OF CANADA
L'ASSOCIATION D'HYPERTENSION
PULMONAIRE DU CANADA



To learn more about pulmonary arterial hypertension (PAH) and available treatments, please visit www.PHACanada.ca/learnmore.

This educational resource was developed by the Canadian Pulmonary Hypertension Professionals Network (CPHPN), a PHA Canada network that brings together PH nurses from across the country to enhance nursing care in the field through leadership, education, and professional development.

The information provided in this resource is for educational purposes only. Please speak with your medical team to establish a personalized treatment plan and discuss any changes to your lifestyle or course of treatment.